

DIMENSION-RAISING THEOREMS FOR COHOMOLOGICAL AND EXTENSION DIMENSIONS

GENCHO SKORDEV AND VESKO VALOV

ABSTRACT. We establish cohomological and extension dimension versions of the Hurewicz dimension-raising theorem

1. INTRODUCTION

Consider a finite-to-one closed and surjective map $f: X \rightarrow Y$. The multiplicity function $\mu: Y \rightarrow \mathbb{N}$ of the map f is defined by $\mu(y) = \text{card}(f^{-1}(y))$, and let $\mu(f) = \{\mu(y) : y \in Y\}$. W. Hurewicz [18] proved that $\text{card}(\mu(f)) \geq k + 1$ provided $\dim Y \geq \dim X + k$ with X and Y being separable metric spaces. Moreover, if $\mu(f) = \{m(1) < m(2) < \dots < m(\ell)\}$, where $\ell \geq k + 1$, then $\dim Y_{m(s+1)}(f) \geq \dim Y - s$ for $s = 0, 1, \dots, k$. Here $Y_m(f) = \{y : \mu(y) = \text{card}(f^{-1}(y)) \geq m\}$. The last assertion is a generalization of the following result of H. Freudenthal [13]: $\dim Y_{s+1}(f) \geq \dim Y - s$, $s = 0, \dots, k$, if $\dim Y \geq \dim X + k$. Hence, $Y_{k+1}(f)$ is not empty which was established first in [17]. The theorem of W. Hurewicz was further generalized for metric spaces in [19, 30], see also [14, 29, 31, 25, 11] for related results.

The present paper deals with cohomological and extension dimension versions of the Hurewicz theorem. Recall that the cohomological dimension $\dim_G$ of the paracompact space X with coefficients in an Abelian group G is defined as follows: $\dim_G X \leq n$ if and only if the Čech cohomology $H^{n+1}(X, A, G) = 0$ for every closed set A in X . Equivalently, $\dim_G X \leq n$ if and only if the homomorphism $H^n(X, G) \rightarrow H^n(A, G)$ of the Čech cohomology groups induced by the inclusion $A \subset X$ is an epimorphism for every closed A in X , see [20, 3, 7]. If Z is a subset of a space X , we define $\text{rdim}_{X,G} Z = \max\{\dim_G F : F \subset Z \text{ is closed in } X\}$.

There exists also another way to define $\dim_G$ for paracompact spaces. If L is a CW -complex, we say that a space X belongs to the class $\alpha(K)$

1991 *Mathematics Subject Classification.* Primary 55M10; Secondary 54F45.

Key words and phrases. metric spaces, cohomological dimension, extension dimension, sheaves.

The second author was partially supported by NSERC Grant 261914-03.

if every map $g: A \rightarrow K$, where $A \subset X$ is closed, can be extended to the whole space X provided g is extendable to some open neighborhood of A in X . Shvedov [27] has shown that if X is paracompact, then $\dim_G X \leq n$ if and only if $X \in \alpha(K(G, n))$, where $K(G, n)$ denotes the Eilenberg-MacLane complex of G in dimension n (for countable groups G this was established by Huber [16]).

Concerning the definition of extension dimension for paracompact spaces, we adopt Dydak's approach [8]. The : If X is a paracompact space and L a CW -complex, we write $e - \dim X \leq L$ if every map $f: A \rightarrow L$, where $A \subset X$ is closed, extends over X up to homotopy. In such a case we say that L is an absolute extensor of X up to homotopy. Let us mention that, according to [8, Proposition 3.5], for every CW -complex L there exists a complete metrizable simplicial complex $K(L)$ homotopy equivalent to L such that, for any paracompact space X we have $K(L)$ is an absolute extensor of X if and only if L is an absolute extensor of X up to homotopy. In case X is compact or metrizable, the above definition of $e - \dim$ coincides with the original one introduced by Dranishnikov [4], i.e., every map $f: A \rightarrow L$ with $A \subset X$ being closed admits an extension over X . The notation $e - \dim X \leq e - \dim Y$ means that $e - \dim X \leq L$ for every CW -complex L with $e - \dim Y \leq L$.

The paper is organized as follows. A cohomological version of the Hurewicz theorem is established in Section 2. Section 3 is devoted to a cohomological version of the mentioned above Freudenthal's theorem. In the final Section 4 we deal with dimension-raising theorems for extension dimension. The first result in this section is an extension analogue of another dimension-raising theorem of Hurewicz stating that if $f: X \rightarrow Y$ is a closed map with $\text{card}(f^{-1}(y)) \leq n + 1$ for every $y \in Y$, then $\dim Y \leq \dim X + n$. This theorem, established first by Hurewicz [17] for separable metric spaces, has many generalizations. For the dimension $\dim$ and normal spaces it was proved by Zarelua [32], Filippov [14] and Pasynkov [23]. Kuzminov [20, Theorem 14, 1), p. 24] was who first provided a cohomological version of this theorem for closed maps between finite dimensional metrizable compact spaces and cohomological dimension $\dim_G$ with respect to an Abelian group G . Kuzminov's proof was based on test spaces for $\dim_G$. With the same method he also established this theorem in the class of paracompact spaces with G being either a periodic group or a field [20, p.39]. Zarelua [32] introduced a new technique for investigating closed finite-to-one maps of paracompact spaces. As a byproduct, he obtained another generalization of the Hurewicz theorem for paracompact spaces and $\dim_G$, where G is a commutative ring with a unity. Finally, for arbitrary Abelian

groups G , paracompact spaces, and more general maps, this theorem was obtained by the first author [31, Corollary 4.1].

The introduction of extension dimension as a unification of the cohomological dimension $\dim_G$ and the ordinary dimension $\dim$ gave another way of possible generalizations of the Hurewicz results. Such a generalization was established by Dranishnikov and Uspenskij [6, Theorem 1.6] for metrizable compacta. Our Proposition 4.1 and Corollary 4.2 extend the Dranishnikov-Uspenskij theorem for more general spaces. Let us mention that, in case of finite-dimensional compacta (resp., metrizable spaces) and simply connected CW -complexes, Proposition 4.1 and Corollary 4.2 follow from the first author's result mentioned above [31, Corollary 4.1] combined with Dranishnikov's homological criterion for extensivity [4, Theorem 9] (resp., with Dydak's criterion [10, Theorems H and G]). We also provide an extension version of the Hurewicz theorem when $\mu(f)$ has finitely many values.

Everywhere below, by a group we mean an Abelian group. Recall that a map $f: X \rightarrow Y$ is perfect if f is a closed map having compact fibers $f^{-1}(y)$, $y \in Y$. It is well known that this is equivalent to f being closed and $f^{-1}(K)$ compact for all compact sets $K \subset Y$.

2. HUREWICZ THEOREM FOR THE COHOMOLOGICAL DIMENSION

Here, we shall prove a theorem for maps increasing the cohomological dimensions $\dim_G$ with coefficients in a group G .

Proposition 2.1. *If $f: X \rightarrow Y$ is a finite-to-one closed surjection between the metric space X and Y and $\dim_G X + k \leq \dim_G Y < \infty$, then $\text{card}(\mu(f)) \geq k + 1$. Moreover, if $\mu(f) = \{m(1) < m(2) < \dots < m(\ell)\}$, then $\ell \geq k + 1$, and $\dim_G Y_{m(s+1)}(f) \geq \dim_G Y - s$ for $s = 0, 1, \dots, k$.*

If $\dim_G X < \dim_G Y = \infty$, then $\text{card}(\mu(f)) \geq \infty$ and $\dim_G Y_m(f) = \infty$ for $m = 2, 3, \dots$

Corollary 2.2. *Let $f: X \rightarrow Y$ be a finite-to-one closed surjection between metrizable spaces and $\text{card}(\mu(f)) \leq k + 1$. Then $\dim_G Y \leq \dim_G X + k$.*

Remarks. 1. *Let $f: X \rightarrow Y$ be a closed surjection between the metrizable spaces X and Y . Then the set $Y_m(f) = \{y : \mu(y) = \text{card}(f^{-1}(y)) \geq m\}$ is a F_σ set in Y , see [11, Lemma 4.3.5, p. 243]. Therefore, $\dim_G Y_m(f) = \text{rdim}_{Y,G} Y_m(f)$.*

2. *Proposition 2.1 and Corollary 2.2 also hold in the more general situation when X is paracompact and submetrizable (recall that a space is*

submetrizable if it admits a continuous metric, or equivalently, it admits a bijective continuous map onto a metrizable space). For the class of submetrizable spaces and the ordinary dimension $\dim$, Proposition 2.1 and Corollary 2.2 were established in [30] (in this case the set $Y_m(f)$ is also F_σ in Y). Bredon [2, Theorem 8.15, p. 243] also provided a version of Corollary 2.2 for separable metric spaces X, Y and a principal ideal domain G .

The proof Proposition 2.1 is based on several lemmas. Before starting the proof, let us provide some notions and results from cohomology and sheaf theory, see [15, 2, 28], and some constructions and theorems of A.V. Zarelua [32, 33, 34]. Recall that the cohomological dimension $\dim_{\mathcal{L}} Z$ of a space Z with coefficients in a sheaf of Abelian groups $\mathcal{L}$ is defined by $\dim_{\mathcal{L}} Z = \min\{n : H^{n+1}(Z, \mathcal{L}_U) = 0 \text{ for every open set } U \subset Z\}$. Moreover, $\dim_{\mathcal{L}} Z = \dim_G Z$ if $\mathcal{L}$ is the constant sheaf $Z \times G$.

First, we recall the notion of a local system of sheaves of groups on a given space Y , see [32, 33, 34]. Let Λ be a partially ordered set. An open covering $\Omega = \{U_\lambda : \lambda \in \Lambda\}$ of Y is said to be Y -directed [34, Definition 7] provided $U_\mu \subset U_\lambda$ for $\lambda \leq \mu$ and every index set $\Lambda_y = \{\lambda \in \Lambda : y \in U_\lambda\}$, $y \in Y$, is directed, i.e., if $y \in U_\lambda \cap U_\mu$, then there exists $\nu \in \Lambda$ such that $\nu \geq \lambda$, $\nu \geq \mu$, and $y \in U_\nu \subset U_\lambda \cap U_\mu$.

Let $\Omega = \{U_\lambda : \lambda \in \Lambda\}$ be an Y -directed open covering. A local system of sheaves $\Sigma = \{\Omega, \mathcal{L}_\lambda, \gamma_\mu^\lambda, \Lambda\}$ is a family of sheaves $\mathcal{L}_\lambda$ on U_λ and homomorphisms $\gamma_\mu^\lambda : \mathcal{L}_\lambda|_{U_\mu} \rightarrow \mathcal{L}_\mu$ with $\mu \geq \lambda$ such that $\nu \geq \mu \geq \lambda$ implies $\gamma_\nu^\lambda = \gamma_\nu^\mu \gamma_\mu^\lambda$, see [34, Definition 13]. A limit $\limind \Sigma$ of the local system of sheaves Σ is defined. This is a sheaf $\mathcal{L}$ on Y with fibers $\mathcal{L}_y = \limind \{(\mathcal{L}_\lambda)_y, \lambda \in \Lambda_y\}$ and the topology of $\mathcal{L}$ is induced by the topology of $\mathcal{L}_\lambda$, [34, Definition 14].

We assume also that the homomorphisms γ_μ^λ are monomorphism, and that for every $U_{\lambda_1}, U_{\lambda_2} \in \Omega$ with $U_{\lambda_1} \cap U_{\lambda_2} \neq \emptyset$ there is $\mu \in \Lambda$ such that $\mu \geq \lambda_1, \lambda_2$ and $U_\mu = U_{\lambda_1} \cap U_{\lambda_2}$. The local inductive systems of sheaves satisfying this condition are called regular, [33, Definition 3.2].

A local system of sheaves on Y , may be considered as a collection of inductive systems of sheaves parameterized by the points of the spaces Y . Then the limit of local system of sheaves may be interpreted as a collection of inductive limits of groups parameterized by the points of the space Y .

Lemma 2.3. [32, Proposition 3.7] *Let $\Sigma = \{\Omega, \mathcal{L}_\lambda, \gamma_\mu^\lambda, \Lambda\}$ be a local system of sheaves on the space Y . If $\text{rdim}(Y, \mathcal{L}_\lambda) \leq n$, then $\dim_{\mathcal{L}} Y \leq n$, where $\mathcal{L} = \limind \Sigma$.*

Here, $\text{rdim}(Y, \mathcal{L}_\lambda) = \max\{\dim_{\mathcal{L}_\lambda} F : F \subset U_\lambda \text{ is a closed set in } Y\}$. More generally, for a fixed open subset U of Y and a sheaf $\mathcal{M}$ on U , $\text{rdim}(Y, \mathcal{M})$ denotes $\max\{\dim_{\mathcal{M}} F : F \subset U, F \text{ is closed in } Y\}$.

If $f: X \rightarrow Y$ is a closed surjective map and G a group, then there is an exact sequence

$$(1) \quad 0 \rightarrow \mathcal{G} \rightarrow f_* f^* \mathcal{G} \rightarrow \mathcal{A} \rightarrow 0,$$

where $\mathcal{G} = Y \times G$, and f_*, f^* are the functors of direct and the inverse image of the sheaves with respect to the map f , see [2], Section I.3. Observe also that $f^* \mathcal{G} = X \times G$.

Remind that a surjective map $f: X \rightarrow Y$ is called zero-dimensional if $\dim f^{-1}(y) = 0$ for every $y \in Y$. This is equivalent to $\dim_G f^{-1}(y) = 0$ for every $y \in Y$ and some (arbitrary) group G .

For closed zero-dimensional maps the sheaf $\mathcal{A}$ is concentrated on the set $Y_2(f)$, i.e., $\mathcal{A}_y = 0$ for $y \notin Y_2(f)$. This fact was used in [29] and will be also exploited here.

Lemma 2.4. *Let $f: X \rightarrow Y$ be a perfect 0-dimensional surjection on the paracompact space X . If $\dim_G X < \dim_G Y < \infty$, then $\dim_{\mathcal{A}} Y = \dim_G Y - 1$, where $\mathcal{A}$ is the sheaf from the exact sequence (1). If $\dim_G X < \dim_G Y = \infty$, then $\dim_{\mathcal{A}} Y = \infty$.*

Proof. Since the map f is closed and 0-dimensional, f_* is an exact functor, see [32, Proposition 1.4]. Then, by [32, Proposition 1.5], $H^i(X, \mathcal{G}_V) = H^i(Y, f_* f^* \mathcal{G}_U)$, where $V = f^{-1}(U)$, $\mathcal{G}_V = (X \times G)_V = f^* \mathcal{G}_U$ and U is an open set in Y . Therefore, $\dim_{f_* f^* \mathcal{G}} Y = \dim_G X$.

First, consider the case $\dim_G X < \dim_G Y < \infty$. The exact sequence (1) implies $\dim_{\mathcal{A}} Y \leq \dim_G Y - 1$. If $m = \dim_G Y$, we can find an open set U in the space Y such that $H^m(Y, \mathcal{G}_U) \neq 0$. Then, according to (1), we have $H^{m-1}(Y, \mathcal{A}_U) \neq 0$ because $m > \dim_G Y$. Hence, $\dim_{\mathcal{A}} Y \geq m - 1$.

Now consider the case $\dim_G X < \dim_G Y = \infty$. Here, $\dim_G Y = \infty$ means that there is a sequence of natural numbers $n_1 < n_2 < \dots$ and open in Y sets U_k such that $H^{n_k}(Y, \mathcal{G}_{U_k}) \neq 0$ for $k = 1, 2, \dots$. Since $H^{n_k}(Y, f_* f^* \mathcal{G}_{U_k}) = H^{n_k}(X, \mathcal{G}_{V_k}) = 0$ for $V_k = f^{-1}(U_k)$, the exact cohomology sequence induced by (1) implies

$$H^{n_k-1}(Y, \mathcal{A}_{U_k}) \rightarrow H^{n_k}(Y, \mathcal{G}_{U_k}) \rightarrow 0$$

for every $n_k > \dim_G X$. Therefore, $H^{n_k-1}(Y, \mathcal{A}_{U_k}) \neq 0$ for all these n_k , which yields $\dim_{\mathcal{A}} Y = \infty$. $\square$

Now, we describe a construction of A. V. Zarelua from his proof of [33, Proposition 3.6]. This construction is given for closed finite-to-one

and surjective maps $f: X \rightarrow Y$ of paracompact spaces X and Y , but it also works for closed zero-dimensional surjective maps.

Local systems $\Sigma_{\mathcal{L}}, \Sigma_{\mathcal{M}}, \Sigma_{\tilde{\mathcal{B}}}$ of sheaves of groups on a space Y are defined as follows. Let $\sigma = \{U_1, \dots, U_{k(\sigma)}\}$ be a disjoint system of open sets in X such that $f^{-1}(U) = U_1 \cup \dots \cup U_{k(\sigma)}$ for some open set U in Y . Consider the set $\Lambda = \{(U, \sigma)\}$ of all pairs (U, σ) satisfying the above conditions and introduce a partial order on Λ determined by the inclusion: $(U', \sigma') \geq (U, \sigma)$ if $U' \subset U$ and σ' is a subdivision of σ . Then $\Omega = \{U_\lambda = U : \lambda = (U, \sigma) \in \Lambda\}$ is an Y -directed open covering.

Let $(U, \sigma) \in \Lambda$, $\sigma = \{U_1, \dots, U_{k(\sigma)}\}$ and

$$\Phi_j = \{x \in U_j : f^{-1}f(x) \setminus U_j \neq \emptyset\}, j = 1, \dots, k(\sigma).$$

Each set Φ_j is closed in U_j . Denote $F_j = f(\Phi_j)$ and let $F = F_1 \cup \dots \cup F_{k(\sigma)}$. Obviously, all sets F and F_j , $j = 1, \dots, k(\sigma)$, are closed in U . Moreover, $F_j \subset Y_2(f)$, $j = 1, \dots, k(\sigma)$.

Consider the sheaves $\mathcal{L}(U, \sigma) = \mathcal{G}_F$, $\mathcal{M}(U, \sigma) = \bigoplus_1^{k(\sigma)} \mathcal{G}_{F_j}$ with $\mathcal{G}$ being the constant sheaf $U \times G$. The natural projections $\mathcal{G}_F \rightarrow \mathcal{G}_{F_j}$ induce a monomorphism $\tilde{\alpha}(U, \sigma) : \mathcal{G}_F \rightarrow \bigoplus_1^{k(\sigma)} \mathcal{G}_{F_j}$. The sheaf $\tilde{\mathcal{B}}(U, \sigma)$ is defined by the exact sequence

$$(2) \quad 0 \rightarrow \mathcal{G}_F \rightarrow \bigoplus_1^{k(\sigma)} \mathcal{G}_{F_j} \rightarrow \tilde{\mathcal{B}}(U, \sigma) \rightarrow 0,$$

i.e., $\tilde{\mathcal{B}}(U, \sigma) = \text{Coker}(\tilde{\alpha}(U, \sigma))$.

If $(U, \sigma), (U', \sigma') \in \Lambda$ with $(U, \sigma) \leq (U', \sigma')$, there are natural homomorphisms

$$\gamma_{(U', \sigma'), \mathcal{L}}^{(U, \sigma)} : \mathcal{L}(U, \sigma)|_{U'} \rightarrow \mathcal{L}(U', \sigma')$$

and

$$\gamma_{(U', \sigma'), \mathcal{M}}^{(U, \sigma)} : \mathcal{M}(U, \sigma)|_{U'} \rightarrow \mathcal{M}(U', \sigma').$$

The last two homomorphisms induce another one

$$\gamma_{(U', \sigma'), \tilde{\mathcal{B}}}^{(U, \sigma)} : \tilde{\mathcal{B}}(U, \sigma)|_{U'} \rightarrow \tilde{\mathcal{B}}(U, \sigma).$$

Then

$$\Sigma_{\mathcal{L}} = \{\Omega, \mathcal{L}(U, \sigma), \gamma_{(U', \sigma'), \mathcal{L}}^{(U, \sigma)}, \Lambda\},$$

$$\Sigma_{\mathcal{M}} = \{\Omega, \mathcal{M}(U, \sigma), \gamma_{(U', \sigma'), \mathcal{M}}^{(U, \sigma)}, \Lambda\}$$

and

$$\Sigma_{\tilde{\mathcal{B}}} = \{\tilde{\mathcal{B}}(U, \sigma), \gamma_{(U', \sigma'), \tilde{\mathcal{B}}}^{(U, \sigma)}, \Lambda\}$$

are local inductive systems of sheaves on the space Y .

Consider also the local inductive system

$$\Sigma_{\tilde{\mathcal{D}}} = \{\tilde{\mathcal{D}}(U, \sigma), \gamma_{(U', \sigma'), \tilde{\mathcal{D}}}^{(U, \sigma)}, \Lambda\},$$

where $\tilde{\mathcal{D}}(U, \sigma) = \bigoplus_1^{k(\sigma)} \mathcal{G}_{U_j}$ and the homomorphisms $\gamma_{(U', \sigma'), \tilde{\mathcal{D}}}^{(U, \sigma)}$ are defined by natural projections.

Let $\mathcal{L} = \limind \Sigma_{\mathcal{L}}$, $\mathcal{M} = \limind \Sigma_{\mathcal{M}}$ and $\tilde{\mathcal{B}} = \limind \Sigma_{\tilde{\mathcal{B}}}$.

Lemma 2.5. [33, Proof of Proposition 3.6] *Let $f : X \rightarrow Y$ be a perfect zero-dimensional surjection between paracompact spaces. Then the sheaves $\mathcal{A}$ and $\tilde{\mathcal{B}}$ are isomorphic.*

Proof. In the case f is finite-to-one, the lemma was established by A.V. Zarelua in [33, Proof of Proposition 3.6] after the construction of the local inductive systems $\Sigma_{\mathcal{L}}$, $\Sigma_{\mathcal{M}}$, $\Sigma_{\tilde{\mathcal{B}}}$ and their limits. The crucial point in that proof, where the assumption that f is finite-to-one is used, is to show that the sheafs $\limind \Sigma_{\tilde{\mathcal{D}}}$ and $f_* f^* \mathcal{G}$ are isomorphic. According to [31, Proposition 1.4], the last fact also holds when f is a zero-dimensional map with all fibers $f^{-1}(y)$, $y \in Y$, being compact. The remaining part of the Zarelua arguments work in the present situation. $\square$

Lemma 2.6. *If $f : X \rightarrow Y$ be a perfect 0-dimensional surjection and X a paracompact space, then $\dim_{\mathcal{A}} Y \leq rdim_{Y,G} Y_2(f)$.*

Proof. The non-trivial case is $rdim_{Y,G} Y_2(f) < \infty$. According to Lemma 2.3, Lemma 2.5 and the exact sequence (2), it suffices to prove the inequalities $rdim(Y, \mathcal{L}(U, \sigma)) \leq rdim_{Y,G} Y_2(f)$ and $rdim(Y, \mathcal{M}(U, \sigma)) \leq rdim_{Y,G} Y_2(f)$ for any $(U, \sigma) \in \Lambda$. On the other hand, since $F = F_1 \cup \dots \cup F_{k(\sigma)}$ and the sets F , F_j , $j = 1, \dots, k(\sigma)$, are closed in U , the second inequality implies the first one. So, we need to prove only the second inequality.

Let C be a closed in Y subset of U . Then $\dim_{\mathcal{G}_{F_i}} C = \dim_{\mathcal{G}_{F_i \cap C}} C = \dim_G C \cap F_i$. Finally, since $C \cap F_i$ is a closed in Y subset of $Y_2(f)$, we have $\dim_G C \cap F_i \leq rdim_{Y,G} Y_2(f)$. This completes the proof. $\square$

Corollary 2.7. *Let X be a paracompact space and $f : X \rightarrow Y$ a perfect 0-dimensional surjection. If $\dim_G X < \dim_G Y < \infty$, then $rdim_{Y,G} Y_2(f) \geq \dim_G Y - 1$. Moreover, if $\dim_G X < \dim_G Y = \infty$, then $rdim_{Y,G} Y_2(f) = \infty$.*

Proof. The proof follows from Lemma 2.4 and Lemma 2.6. $\square$

Remark. *The second part of Corollary 2.7 also holds when f is a closed zero-dimensional surjection (not necessarily perfect), but X and Y being metrizable. Indeed, then $\dim_{Y,G} Y_2(f) = \dim_G Y_2(f)$ and assume that $\dim_G Y_2(f) < \infty$. Let $Y^1 = Y \setminus Y_2(f)$ and $X^1 = f^{-1}Y^1$. The map $f_1 = f|_{X^1} : X^1 \rightarrow Y^1$ is a homeomorphism. Therefore, $\dim_G Y^1 = \dim_G X^1$. Moreover, $\dim_G X^1 \leq \dim_G X$, [20, Theorem 16,*

2]. Hence, by [9, Theorem B], $\dim_G Y \leq \dim_G Y^1 + \dim_G Y_2(f) + 2 < \infty$, which is a contradiction.

Lemma 2.8. *Let $f : X \rightarrow Y$ be a closed 0-dimensional surjection between metrizable spaces and $m \geq 2$. If $\dim_G X < \dim_G Y_m(f) < \infty$, then $\dim_{Y,G} Y_{m+1}(f) \geq \dim_{Y,G} Y_m(f) - 1$. If $\dim_G Y_m(f) = \infty$, then $\dim_G Y_{m+1}(f) = \infty$.*

Proof. The set $X_1 = \cup\{Bd(f^{-1}(y)) : y \in Y\}$, where $Bd(f^{-1}(y))$ is the boundary of $f^{-1}(y)$ in X , is closed in X because the map f is closed, [22, Lemma 23-2, p. 142]. So, $f_1 = f|_{X_1} : X_1 \rightarrow Y$ is also closed, surjective and zero-dimensional. By the Veinstein's lemma [11, 1.12.9, p. 111], all sets $f_1^{-1}(y)$, $y \in Y$, are compact. Moreover, $\dim_G X_1 \leq \dim_G X$. Therefore, without restriction of the generality, we can assume that the original map f is perfect.

First consider the case $\dim_G X < \dim_G Y_m(f) < \infty$. Choose locally finite closed coverings $\omega^j = \{F_s^j\}_s$ of X with $\text{diam} F_s^j \leq 1/j$, $j = 1, 2, \dots$. Since $Y_m(f)$ is an F_σ -set in Y , there is a closed set $C \subset Y$ with $C \subset Y_m(f)$ and $\dim_G C = \dim_G Y_m(f)$. Let $\tilde{C} = f^{-1}(C)$ and $f_C = f|_{\tilde{C}} : \tilde{C} \rightarrow C$. The map f_C is perfect, zero-dimensional and surjective. Moreover,

$$(3) \quad Y_m(f_C) = \{y \in C : \text{card}(f_C^{-1}(y)) \geq m\} = C.$$

Let $y \in C$ and $\{x_1, \dots, x_m\} \subset f^{-1}(y) = f_C^{-1}(y)$. There exists $j(y)$ such that the covering $\omega^{j(y)}$ separates the points $\{x_1, \dots, x_m\}$, i.e., we can find disjoint sets $F_{s_1}^{j(y)}, \dots, F_{s_m}^{j(y)} \in \omega^{j(y)}$ such that $x_i \in F_{s_i}^{j(y)}$, $i = 1, \dots, m$.

Consider the closed subsets $F_{j(y)} = \cap_{i=1}^m f_C(F_{s_i}^{j(y)} \cap \tilde{C})$ of C and let $C_j = \cup_{j(y)=j} F_{j(y)}$. Since f is closed and the fibers $f^{-1}(y)$, $y \in Y$, are compact, all systems $\{F_{j(y)}\}_{j(y)=j}$, $y \in C$, are locally-finite and consist of closed sets. Therefore, each set C_j is closed in C . Moreover, (3) implies $C = \cup_j C_j$. Then, there exist j_0 and $y_0 \in C$ with $j_0 = j(y_0)$ and $\dim_G Y_m(f) = \dim_G C = \dim_G C_{j_0} = \dim_G F_{j_0}$. The sets $F_{s_1}^{j_0}, \dots, F_{s_m}^{j_0} \in \omega^{j_0}$ separate the points $\{x_1, \dots, x_m\} \subset f^{-1}(y_0)$. Let $\Phi = F_{s_1}^{j_0} \cap f^{-1}(F_{j_0})$ and $f_\Phi = f|_\Phi : \Phi \rightarrow F_{j_0}$. The map f_Φ is perfect, surjective and 0-dimensional and we have the following inclusion

$$(4) \quad Y_2(f_\Phi) = \{y \in F_{j_0} : \text{card}(f_\Phi^{-1}(y)) \geq 2\} \subset Y_{m+1}(f).$$

Moreover, $\dim_G \Phi \leq \dim_G X < \dim_G Y_m(f) = \dim_G F_{j_0} < \infty$. Then, by Corollary 2.2, $\dim_G Y_2(f_\Phi) \geq \dim_G F_{j_0} - 1 = \dim_G Y_m(f) - 1$. Finally, (4) completes the proof in the case $\dim_G Y_m(f) < \infty$.

Now consider the case $\dim_G Y_m(f) = \infty$. Since $Y_m(f)$ is a F_σ set in Y , we have two cases: (A) There is a sequence of natural numbers

$n_1 < n_2 < \dots$ converging to ∞ and closed in Y sets $C^k \subset Y_m(f)$ such that $\dim_G C^k = n_k$ for $k = 1, 2, \dots$; (B) There is a closed in Y set $C \subset Y_m(f)$ such that $\dim_G C = \infty$.

Case (A). Let $n_k > \dim_G X$. Applying the construction from the proof of the lemma in the case $\dim_G Y(f) < \infty$ (with C replaced by C^k), we obtain $\dim_G Y_{m+1} \geq n_k - 1$. This implies $\dim_G Y_{m+1} = \infty$.

Case (B). We apply again the construction from the case $\dim_G Y_m(f) < \infty$ for the set C to obtain $C = \cup_j C_j$, where $C_j = \cup_{j(y)=j} F_{j(y)}$, $y \in C$. Since $\dim_G C = \infty$, we have again two cases: (B₁) There exists a sequence of natural numbers $n_1 < n_2 < \dots$ converging to ∞ and closed sets $C_{m_k} \subset C$ such that $\dim_G C_{m_k} = n_k$; (B₂) There exists j_0 such that $\dim_G C_{j_0} = \infty$ for some closed $C_{j_0} \subset C$.

In the Case (B₁) we argue as in the Case (A) and obtain $\dim_G Y_{m+1} = \infty$.

The Case (B₂) splits again in two cases: (B₂₁) There is a sequence of natural numbers $n_1 < n_2 < \dots$ such that $\lim n_k = \infty$ and $\dim_G F_{j(y_k)} = n_k$ for $y_k \in C_{j_0}$ and $j(y_k) = j_0$; (B₂₂) There exists $y_0 \in C_{j_0}$ such that $\dim_G F_{j(y_0)} = \infty$.

The arguments from Case (A) applied under the hypotheses of Case (B₂₁) imply $\dim_G Y_{m+1} = \infty$.

In Case (B₂₂) we apply the construction from the case $\dim_G Y_m(f) < \infty$ for the set C and consider the map $f_\Phi : \Phi \rightarrow F_{j(y_0)}$. This map is perfect, zero-dimensional and surjective. Moreover, $\dim_G \Phi \leq \dim_G X$ and $\dim_G F_{j(y_0)} = \infty$. Then, by Corollary 2.7, $\dim_G Y_2(f_\Phi) = \infty$. Finally, inclusion (4) yields $\dim_G Y_{m+1} = \infty$. $\square$

Proof of Proposition 2.1. Assume $\dim_G X + k \leq \dim_G Y < \infty$, $k \geq 1$, and $\mu(f) = \{m_1 < \dots < m_\ell\}$. Then, by Corollary 2.7 and Lemma 2.8,

$$\dim_G Y_{m_{k+1}}(f) \geq \dim_G X.$$

So, $Y_{m_{k+1}}(f)$ is not empty, which yields $\ell \geq k + 1$.

If $\dim_G X < \dim_G Y = \infty$, again by Corollary 2.7 and Lemma 2.8, we have $\dim_G Y_m(f) = \infty$ for $m \geq 2$. Therefore, $\text{card}(\mu(f)) = \infty$.

Another result of Hurewicz [18, Theorem II, p.74-76] is stating that if $f : X \rightarrow Y$ a closed surjective finite-to-one map between separable metrizable spaces, X has the property (α) and $\dim Y - \dim X \geq n$, then $\text{card}(\mu(f)) \geq n + 2$.

Here, we say that a space X has the property (α) if $\dim A < \dim X$ for every nowhere dense closed set A in X . For example, every topological manifold X satisfies the property (α) . In his proof Hurewicz used that the space Y can be assumed to satisfy the following condition (β) : $\dim U = \dim Y$ for every open set U in Y .

A version of this theorem also holds for the cohomological dimension. We say that $X \in (\alpha)_G$, where G is a group, if $\dim_G A < \dim_G X$ for every nowhere dense closed set $A \subset X$. Let us remind that every locally compact n -cohomological manifold M over the principal ideal domain L , notation M is $n - cm_L$, possesses the property $(\alpha)_L$, see [1, Proposition 4.9(a), p.14]. We also consider the corresponding analogue of condition (β) : A space Y satisfies condition $(\beta)_G$, notation $Y \in (\beta)_G$, if $\dim_G U = \dim_G Y$ for every open $U \subset Y$.

Now, we can state a version of the Hurewicz result mentioned above.

Proposition 2.9. *Let $f : X \rightarrow Y$ be a closed, surjective and finite-to-one map on the metrizable space X . If $X \in (\alpha)_G$, $\dim_G X + n \leq \dim_G Y < \infty$ and $Y \in (\beta)_G$, then $\text{card}(\mu(f)) \geq n + 2$.*

W. Hurewicz derived his theorem from a lemma whose cohomological version is Lemma 2.10 below. Observe that Proposition 2.9 follows from Lemma 2.10 and the inequality (5).

Lemma 2.10. *Let $f : X \rightarrow Y$ be a closed finite-to-one map between the metrizable spaces X and Y . Assume that $X \in (\alpha)_G$ and $Y \in (\beta)_G$. If $\dim_G X + k \leq \dim_G Y < \infty$ and $m_0 = \max\{\mu(y) : y \in Y\}$, then $\dim_G Y_{m_0} < \dim_G X$.*

Proof. The proof of this lemma adapts the arguments from the proof of the corresponding assertion for the dimension $\dim$, [18, p.75-76]. Hurewicz worked with a countable basis of the space X . Instead of this, we take the collection of elements of the closed coverings ω^j , see the proof of Lemma 2.8, Case (A). Then we can apply the constructions of W. Hurewicz with $\dim$ replaced by $\dim_G$. $\square$

3. FREUDENTHAL'S THEOREM FOR THE COHOMOLOGICAL DIMENSION

In this section we shall prove the following

Proposition 3.1. *Let $f : X \rightarrow Y$ be a perfect 0-dimensional surjection between the paracompact spaces X and Y . Suppose $\dim_G X + k \leq \dim_G Y < \infty$ for some natural number k and a countable group G . Then $\text{rdim}_{Y,G} Y_{s+1}(f) \geq \dim_G Y - s$ for every $s = 1, \dots, k$.*

First, we need the following technical result.

Lemma 3.2. *Let $f : X \rightarrow Y$ be a 0-dimensional closed and surjective map between the paracompact spaces X and Y and G be a countable group. Suppose F is a closed in Y subset of $Y_2(f)$ and let $\Phi = f^{-1}(F)$. Then there exist families $\{\Phi_{\alpha,g} : \alpha \in \Lambda, g \in G\}$ and $\{F_{\alpha,g} : \alpha \in \Lambda, g \in G\}$ of closed sets in X and Y , respectively, such that:*

- (1) $\Phi = \cup_{\alpha,g} \Phi_{\alpha,g}$, $F = \cup_{\alpha,g} F_{\alpha,g}$ and $f(\Phi_{\alpha,g}) = F_{\alpha,g}$;
- (2) $\{F_{\alpha,g} : \alpha \in \Lambda, g \in G\}$ is locally countable in F and, for every $g \in G$ the family $\{F_{\alpha,g} : \alpha \in \Lambda\}$ is locally finite in F ;
- (3) $Y_t(\varphi_{\alpha,g}) = \{y \in F_{\alpha,g} : \text{card}(\varphi_{\alpha,g}^{-1}(y)) \geq t\} \subset Y_{t+1}(f)$, where $\varphi_{\alpha,g} : \Phi_{\alpha,g} \rightarrow F_{\alpha,g}$ is the restriction map $f|_{\Phi_{\alpha,g}}$.

Proof. Recall that the support of the sheaf $\mathcal{A}$ is the set $Y_2(f)$. For every $y \in Y$ choose an open in Y set O_y and a section $s_y \in \Gamma(\overline{O_y}, \mathcal{A})$ such that $s_y(z) \neq 0$ for all $z \in \overline{O_y} \cap F$. Here $\overline{O_y}$ is the closure of the set O_y in Y . Assume that O_y is so small that there exists a section $\tilde{s}_y \in \Gamma(\overline{O_y}, f_*G)$ such that s_y corresponds to $\tilde{s}_y$ by the natural projection $f_*G \rightarrow \mathcal{A}$ from the exact sequence (1). Let $t_y \in \Gamma(f^{-1}O_y, G)$ be the section which corresponds to the section $\tilde{s}_y$ by the isomorphism of the groups of sections $\Gamma(O_y, f_*G)$ and $\Gamma(f^{-1}O_y, G)$. Consider the covering $\omega = \{O_y \cap F\}_{y \in F}$ and let $\tilde{\omega} = \{F_\alpha : \alpha \in \Lambda\}$ be a closed locally finite covering of F refining ω . Then, for every α there is $y(\alpha) \in F$ such that $F_\alpha \subset \overline{O_{y(\alpha)}}$. Let $s_\alpha = s_{y(\alpha)}$, $\tilde{s}_\alpha = \tilde{s}_{y(\alpha)}$, $t_\alpha = t_{y(\alpha)}$ and $\Phi_\alpha = f^{-1}(F_\alpha)$.

Consider the sets $\Phi_{\alpha,g} = \{x \in \Phi_\alpha : t_\alpha(x) = g\}$, $\alpha \in \Lambda$, $g \in G$. Each $\Phi_{\alpha,g}$ is closed in X because the sheaf $X \times G$ is a Hausdorff space. Moreover, since G is countable, the family $\{\Phi_{\alpha,g} : g \in G\}$ is countable for every $\alpha \in \Lambda$. Obviously, $\Phi_\alpha = \cup_{g \in G} \Phi_{\alpha,g}$, and let $F_{\alpha,g} = f(\Phi_{\alpha,g})$. The family $\{F_{\alpha,g} : \alpha \in \Lambda\}$ is locally finite in F for every $g \in G$ because so is the family $\{F_\alpha : \alpha \in \Lambda\}$. This implies that $\{F_{\alpha,g} : \alpha \in \Lambda, g \in G\}$ is a locally countable family in F . Obviously, every element of the last family is a closed subset of Y .

Denote by $\varphi_{\alpha,g}$ the restriction of f on the set $\Phi_{\alpha,g}$. For every $y \in F_{\alpha,g}$ we have $s_\alpha(y) \neq 0$. Hence, there exist points $x_1, x_2 \in f^{-1}(y)$ such that $t_\alpha(x_1) \neq t_\alpha(x_2)$. Indeed, otherwise $t_\alpha(x') = t_\alpha(x'')$ for all $x', x'' \in f^{-1}(y)$ would imply $s_\alpha(y) = 0$, which is not possible. So, $x_i \in \Phi_{\alpha,g_i}$, where $g_i = t_\alpha(x_i)$, $i = 1, 2$. Therefore,

$$Y_t(\varphi_{\alpha,g}) = \{y \in F_{\alpha,g} : \text{card}(\varphi_{\alpha,g}^{-1}(y)) \geq t\} \subset Y_{t+1}(f).$$

This completes the proof. $\square$

Proof of Proposition 3.1

The proof goes by induction with respect to k . Corollary 2.7 provides the case $k = 1$. Assume the assertion has been proved for all k with $1 \leq k \leq \ell - 1$ and let $k = \ell$.

Choose a closed in Y set F which is contained in $Y_2(f)$ such that $\dim_G F = r \dim_{Y,G} Y_2(f) \geq \dim_G Y - 1 > \dim_G X$. Consider the families $\{\Phi_{\alpha,g} : \alpha \in \Lambda, g \in G\}$ and $\{F_{\alpha,g} : \alpha \in \Lambda, g \in G\}$ from Lemma 3.2. Since $F = \cup_{\alpha,g} F_{\alpha,g}$, G is countable and $\{F_{\alpha,g} : \alpha \in \Lambda\}$ is locally

finite in F for every $g \in G$, $\dim_G F_{\alpha,g} = \dim_G F$ for some $\alpha \in \Lambda$ and $g \in G$. Then both $F_{\alpha,g}$ and $\Phi_{\alpha,g}$ are paracompact spaces and the map $f_{\alpha,g} : \Phi_{\alpha,g} \rightarrow F_{\alpha,g}$ is perfect, surjective and zero-dimensional. Moreover,

$$\dim_G F_{\alpha,g} = \dim_G F \geq \dim_G Y - 1 \geq \dim_G X + k - 1 \geq \dim_G \Phi_{\alpha,g} + k - 1.$$

It follows from the inductive assumption that

$$rdim_{Y,G} Y_s(f_{\alpha,g}) = rdim_{\Phi_{\alpha,g},G} Y_s(f_{\alpha,g}) \geq \dim_G F_{\alpha,g} - s + 1$$

for $s = 1, \dots, k - 1$. According to Lemma 3.2(3), we have $Y_s(f_{\alpha,g}) \subset Y_{s+1}(f)$. Then

$$rdim_{Y,G} Y_{s+1}(f) \geq rdim_{Y,G} Y_s(f_{\alpha,g}) \geq \dim_G Y - s$$

for $s = 1, \dots, k$, which is exactly what we need.

4. HUREWICZ THEOREM FOR EXTENSION DIMENSION

In this section we establish extension dimensional analogues of the Hurewicz theorems when either $\text{card}(f^{-1}(y)) \leq n + 1$ for every $y \in Y$ or $\text{card}(\mu(f)) \leq n + 1$. A map $f : X \rightarrow Y$ is said to have a metrizable kernel if there exist a metrizable space M and a map $g : X \rightarrow M$ such that g is injective on each fiber of f , i.e., $g|_{f^{-1}(y)}$ is an one-to-one map for every $y \in Y$. Obviously, this is the case if X is submetrizable.

Proposition 4.1 below was established by Dranishnikov and Uspenskij [6, Theorem 1.6] in the case both X and Y are metrizable compacta. Our proof of this proposition is a modification of the Dranishnikov and Uspenskij arguments.

Proposition 4.1. *Let $f : X \rightarrow Y$ be a closed surjective map admitting a metrizable kernel such that Y is a paracompact k -space and $\text{card}(f^{-1}(y)) \leq n + 1$ for every $y \in Y$. Then $e - \dim Y \leq e - \dim(X \times \mathbb{I}^n)$.*

Proof. Let $m = n + 1$ and $P_m(X)$ be the space of all probability measures on X whose supports consist of at most m points. The map f has an extension $P_m(f) : P_m(X) \rightarrow P_m(Y)$. Actually, $P_m(f)$ is the restriction of the map $P_m(\beta f) : P_m(\beta X) \rightarrow P_m(\beta Y)$ on $P_m(X)$, where $\beta f : \beta X \rightarrow \beta Y$ is the natural extension of f between the Čech-Stone compactifications of X and Y . Since f is a perfect map, so is $P_m(f)$. Hence, $Z = P_m(f)^{-1}(Y)$ is a paracompact k -space as a perfect preimage of Y . If Δ_n denotes the standard n -dimensional simplex in $\mathbb{R}^n$, there exists a map $p : X^m \times \Delta_n \rightarrow P_m(X)$ assigning to each point $(x_0, x_1, \dots, x_n, t_0, t_1, \dots, t_n) \in X^m \times \Delta_n$ the measure $\mu = \sum_{i=0}^n t_i x_i$. We

consider another map $\pi: T \rightarrow X \times \Delta_n$, where $T = p^{-1}(Z)$, defined by $\pi(x_0, x_1, \dots, x_n, t_0, t_1, \dots, t_n) = (x_0, t_0, t_1, \dots, t_n)$. Obviously, π is surjective.

Claim. The maps $p|_T$ and π are perfect. Moreover T is closed in $X^m \times \Delta_n$.

Since Z is a k -space, according to [12, Theorem 3.7.18], it suffices to show that $p^{-1}(K)$ is a compact subset of T for every compact set $K \subset Z$. So, we fix such K and consider the compact set $F = f^{-1}(P_m(f)(K)) \subset X$. Then $F^m \times \Delta_n$ is a compact subset of $X^m \times \Delta_n$ containing $p^{-1}(K)$. On the other hand $p^{-1}(K)$ is closed in $X^m \times \Delta_n$ because K is closed in $P_m(X)$. Therefore, $p^{-1}(K)$ is compact. Obviously, X is k -space as a closed subset of Z , so is $X \times \Delta_n$. Consequently, to prove that π is perfect, again by [12, Theorem 3.7.18], it suffices to show that $\pi^{-1}(K \times \Delta_n)$ is compact for any compact $K \subset X$. Since $\pi^{-1}(K \times \Delta_n)$ is contained in $K^m \times \Delta_n$, the proof of the claim is reduced to establish that T is closed in $X^m \times \Delta_n$. The last assertion follows from the fact that Z is closed in $P_m(X)$ (because $Z = P_m(f)^{-1}(Y)$ and Y is closed in $P_m(Y)$) and $T = p^{-1}(Z)$.

We are going to show that $e - \dim T \leq e - \dim(X \times \mathbb{I}^n)$. To this end, we need the following result of Dranishnikov and Uspenskij [6, Theorem 1.2]: If $g: Z_1 \rightarrow Z_2$ is a perfect 0-dimensional surjection between paracompact spaces, then $e - \dim Z_1 \leq e - \dim Z_2$ (this theorem was originally established for compact spaces, but the proof works for paracompact spaces as well). Since π is perfect and, obviously, each fiber of π is 0-dimensional, the above theorem yields $e - \dim T \leq e - \dim(X \times \mathbb{I}^n)$.

Next step is to show that $e - \dim Z \leq e - \dim(X \times \mathbb{I}^n)$. We fix a metric space (M, ρ) and a map $g: X \rightarrow M$ such that g is injective on each fiber $f^{-1}(y)$, $y \in Y$. Let d be the continuous pseudometric on X defined by $d(x', x'') = \rho(g(x'), g(x''))$ and consider the corresponding pseudometric d_m on X^m , $d_m((x_i), (y_i)) = \max_{0 \leq i \leq n} d(x_i, y_i)$. For every equivalence relation $\mathfrak{R}$ on the set $\{0, 1, \dots, n\}$ and every $\varepsilon > 0$ let $E_{\mathfrak{R}, \varepsilon} \subset X^m$ be the set of all m -tuples $(x_0, \dots, x_n)$ satisfying the following conditions:

- if $i \mathfrak{R} j$, then $x_i = x_j$;
- if i is not $\mathfrak{R}$ -equivalent to j , then $d(x_i, x_j) \geq \varepsilon$.

Since d is a continuous pseudometric on X , $E_{\mathfrak{R}, \varepsilon}$ is a closed subset of X^m . We choose one element from every equivalence class of $\mathfrak{R}$ and denote by $S_{\mathfrak{R}}$ the set of these $\mathfrak{R}$ -representatives. Let also

$$\Delta_{\mathfrak{R}} = \{t_0, \dots, t_n\} \in \Delta_n : t_i > 0 \text{ if and only if } i \in S_{\mathfrak{R}}\}$$

As in the proof of Theorem 1.7 from [6], one can show that the restriction of p on the set $C \times \Delta_{\mathfrak{R}}$ is one-to-one for every closed set

$C \subset E_{\mathfrak{R},\varepsilon}$ with a d_m -diameter $\leq \varepsilon/2$. Since $\Delta_{\mathfrak{R}}$ is an F_σ -subset of Δ_n , $(C \times \Delta_{\mathfrak{R}}) \cap T$ is F_σ in T .

Now, for every $\varepsilon > 0$ choose a locally finite closed covering $\omega_\varepsilon = \{H_{\alpha,\varepsilon} : \alpha \in \Lambda_\varepsilon\}$ of X^m with each $H_{\alpha,\varepsilon}$ having a d_m -diameter $\leq \varepsilon/2$ (this can be done as follows: first, choose a locally finite closed covering γ_ε of M^m such that the ρ_m -diameter of each element of γ_ε is $\leq \varepsilon/2$, where $\rho_m((a_i), (b_i)) = \max_{0 \leq i \leq n} \rho(a_i, b_i)$, and then let ω_ε to be $(g^m)^{-1}(\gamma_\varepsilon)$). Next, consider the locally finite in T families

$$\Theta_{\mathfrak{R},k} = \{((H_{\alpha,1/k} \cap E_{\mathfrak{R},1/k}) \times \Delta_{\mathfrak{R}}) \cap T : \alpha \in \Lambda_{1/k}\}$$

with $k \in \mathbb{N}$ and $\mathfrak{R}$ being an equivalence relation on $\{0, 1, \dots, n\}$. Each element of $\Theta_{\mathfrak{R},k}$ is an F_σ -subset of T , so $((H_{\alpha,1/k} \cap E_{\mathfrak{R},1/k}) \times \Delta_{\mathfrak{R}}) \cap T = \bigcup_{j=1}^\infty H_{\mathfrak{R},k,j}(\alpha)$ such that all $H_{\mathfrak{R},k,j}(\alpha)$ are closed in T . Therefore, we obtain countably many families

$$\Omega_{\mathfrak{R},k,j} = \{p(H_{\mathfrak{R},k,j}(\alpha)) : \alpha \in \Lambda_{1/k}\}$$

of closed subsets of Z . Moreover, we already observed that p restricted to $H_{\mathfrak{R},k,j}(\alpha)$ is bijective. Because $p|T$ is a closed map, this yields that $H_{\mathfrak{R},k,j}(\alpha)$ and $p(H_{\mathfrak{R},k,j}(\alpha))$ are homeomorphic. So, $e - \dim p(H_{\mathfrak{R},k,j}(\alpha)) \leq e - \dim(X \times \mathbb{I}^n)$. On the other hand, since all $\Theta_{\mathfrak{R},k}$ are locally finite in T and $p|T$ is perfect, the families $\Omega_{\mathfrak{R},k,j}$ are locally finite in Z . Therefore, each set $Z_{\mathfrak{R},k,j} = \bigcup \{p(H_{\mathfrak{R},k,j}(\alpha)) : \alpha \in \Lambda_{1/k}\}$ is closed in Z and, according to the locally finite sum theorem for extension dimension [5, Proposition 1.18], $e - \dim Z_{\mathfrak{R},k,j} \leq e - \dim(X \times \mathbb{I}^n)$ for any $\mathfrak{R}$ and $k, j \in \mathbb{N}$. Since d is a metric on each fiber of f , it is easily seen that Z is the union of the sets $Z_{\mathfrak{R},k,j}$, $k, j \in \mathbb{N}$ and $\mathfrak{R}$ being an equivalence relation on $\{0, 1, \dots, n\}$. Therefore, by the countable sum theorem for extension dimension, $e - \dim Z \leq e - \dim(X \times \mathbb{I}^n)$.

The last step of our proof is to show that $e - \dim Y \leq e - \dim(X \times \mathbb{I}^n)$. We are going to use the following result of Dranishnikov and Uspenskij [6, Proposition 2.3]: Every surjective perfect map $g: Z_1 \rightarrow Z_2$ with convex fibers between paracompact spaces has the following property: for any CW -complex K and every closed subset $B \subset Z_2$ the restriction $g: g^{-1}(B) \rightarrow B$ induced a bijective map $g^*: [B, K] \rightarrow [g^{-1}(B), K]$ between the homotopy classes (such maps are called hereditary shape equivalences). This result was established for compact spaces but its proof holds for paracompact spaces as well. Here, a map $g: Z_1 \rightarrow Z_2$ has convex fibers if there exists a convex subset E of a locally convex linear space and a closed embedding $j: Z_1 \subset Z_2 \times E$ such that the sets $Z_1(y) = \{x \in E : (y, x) \in j(Z_1)\}$ are convex and compact for every $y \in Z_2$. The idea behind the proof of this result is the following simple fact: If $\phi: \bar{Y} \rightarrow \bar{Z}$ is an upper semicontinuous set-valued map with

compact and convex values, where $\bar{Y}$ is paracompact and $\bar{Z}$ is a convex subset of a locally convex linear space, then for every family $\mathcal{U}$ of open in $\bar{Z}$ sets with each $\phi(y)$, $y \in \bar{Y}$, being a subset of some $U \in \mathcal{U}$, there exists a map $h: \bar{Y} \rightarrow \bar{Z}$ such that h and ϕ are $\mathcal{U}$ -close, i.e., for every $y \in \bar{Y}$ both $h(y)$ and $\phi(y)$ are contained in an element of $\mathcal{U}$.

In our situation, we consider the perfect map $P_m(f): Z \rightarrow Y$ which has convex fibers. Indeed, let $P(\beta X)$ be the space of all probability measures on βX . It is well known that $P(\beta X)$ is a compact and convex subset of $\mathbb{R}^{C(\beta X)}$, $C(\beta X)$ being the set of all continuous functions on βX . Moreover, there exists a natural map $P(\beta f): P(\beta X) \rightarrow P(\beta Y)$ extending $P_m(f)$. It is easily seen that the map $j: Z \rightarrow Y \times P(\beta X)$, $j(\mu) = (P(\beta f)(\mu), \mu)$, is a closed embedding and all sets $P_m(f^{-1}(y)) = \{\mu \in P(\beta X) : (y, \mu) \in j(Z)\}$, $y \in Y$, are convex and compact. So, by the mentioned above result [6, Proposition 2.3], $P_m(f)$ is a hereditary shape equivalence. Since $e - \dim Z \leq e - \dim(X \times \mathbb{I}^n)$, this implies the required inequality $e - \dim Y \leq e - \dim(X \times \mathbb{I}^n)$. $\square$

Corollary 4.2. *Let $f: X \rightarrow Y$ be a surjective map between the compact spaces X and Y with $\text{card}(f^{-1}(y)) \leq n + 1$ for every $y \in Y$. Then $e - \dim Y \leq e - \dim(X \times \mathbb{I}^n)$.*

Proof. Suppose L is a CW -complex such that $e - \dim(X \times \mathbb{I}^n) \leq L$ and $m = n + 1$. Obviously, X and Y have the same topological weight. If, $w(X) = w(Y)$ is countable, the proof follows directly from Proposition 4.1. Otherwise, using the notations from the proof of Proposition 4.1, consider the spaces $Z = P_m(f)^{-1}(Y)$, $T = p^{-1}(Z)$ and the diagram $T \xrightarrow{p} Z \xrightarrow{P_m(f)} Y$. According to the Ščepin spectral theorem [26], we can find continuous ω -inverse systems $S_X = \{X_\alpha, \xi_\alpha^\beta : \alpha, \beta \in A\}$ and $S_Y = \{Y_\alpha, \varpi_\alpha^\beta : \alpha, \beta \in A\}$ consisting of metrizable compacta, and continuous maps $f_\alpha: X_\alpha \rightarrow Y_\alpha$, $\alpha \in A$, such that $X = \varprojlim S_X$, $Y = \varprojlim S_Y$ and $\varpi_\alpha \circ f = \xi_\alpha \circ f_\alpha$. Here, $\xi_\alpha: X \rightarrow X_\alpha$ and $\varpi_\alpha: Y \rightarrow Y_\alpha$ are the projections of the inverse systems S_X and S_Y , respectively. Then $P_m(X) = \varprojlim \{P_m(X_\alpha), P_m(\xi_\alpha^\beta) : \alpha, \beta \in A\}$ and $X^m \times \Delta_n = \varprojlim \{X_\alpha^m \times \Delta_n, (\xi_\alpha^\beta)^m \times id : \alpha, \beta \in A\}$. This implies $T = \varprojlim \{T_\alpha, (\xi_\alpha^\beta)^m \times id : \alpha, \beta \in A\}$ and $Z = \varprojlim \{Z_\alpha, P_m(\xi_\alpha^\beta) : \alpha, \beta \in A\}$, where $T_\alpha = (\xi_\alpha^m \times id)(T)$ and $Z_\alpha = P_m(\xi_\alpha)(Z)$. We also consider the maps $p_\alpha: T_\alpha \rightarrow Z_\alpha$ assigning to each point $(\xi_\alpha(x_0), \xi_\alpha(x_1), \dots, \xi_\alpha(x_n), t_0, t_1, \dots, t_n) \in X_\alpha^m \times \Delta_n$ the measure $\sum_{i=0}^n t_i \xi_\alpha(x_i) \in Z_\alpha$. Therefore, the following diagrams are commutative for all α :

$$\begin{array}{ccccccccc}
T & \xrightarrow{p} & Z & \xrightarrow{P_m(f)} & Y & \xleftarrow{f} & X & \xrightarrow{id} & P_m(X) \\
\downarrow \xi_\alpha^m \times id & & \downarrow P_m(\xi_\alpha) & & \downarrow \varpi_\alpha & & \downarrow \xi_\alpha & & \downarrow P_m(\xi_\alpha) \\
T_\alpha & \xrightarrow{p_\alpha} & Z_\alpha & \xrightarrow{P_m(f_\alpha)} & Y_\alpha & \xleftarrow{f_\alpha} & X_\alpha & \xrightarrow{id} & P_m(X_\alpha)
\end{array}$$

Let us observe that the requirement for f in Proposition 4.1 to have a metrizable kernel was necessary only to prove that $e - \dim Z \leq e - \dim T$, the other two facts we established there remain valid in the present situation. So, $e - \dim T \leq e - \dim(X \times \mathbb{I}^n)$ and $e - \dim Y \leq e - \dim Z$. Thus, we need to show that $e - \dim Z \leq e - \dim(X \times \mathbb{I}^n)$.

Since $e - \dim T \leq e - \dim(X \times \mathbb{I}^n) \leq L$, we can use the factorization theorem for extension dimension (see [24, Theorem 2], [21]) to construct the inverse system $S_T = \{Z_\alpha, P_m(\xi_\alpha^\beta) : \alpha, \beta \in A\}$ in such a way that $e - \dim T_\alpha \leq L$, $\alpha \in A$. The equalities $Z_\alpha = P_m(f_\alpha)^{-1}(Y_\alpha)$ and $T_\alpha = p_\alpha^{-1}(Z_\alpha)$ may not be true for some α , but we always have the inclusions $Z_\alpha \subset P_m(f_\alpha)^{-1}(Y_\alpha)$ and $T_\alpha \subset p_\alpha^{-1}(Z_\alpha)$. Since each f_α has a metrizable kernel, according to the arguments from the proof of Proposition 4.1, $e - \dim Z_\alpha \leq e - \dim T_\alpha$. Consequently, $e - \dim Z_\alpha \leq L$ for every α . Finally, Z being the limit of the inverse system S_T yields $e - \dim Z \leq L$. $\square$

Next proposition provides the relation between $e - \dim Y$ and $e - \dim X$ when $f: X \rightarrow Y$ is a finite-to-one closed map with $\text{card}(\mu(f)) \leq n + 1$. For any CW -complexes K and L their join is denoted by $K \star L$.

Proposition 4.3. *Let $f: X \rightarrow Y$ be a closed surjection between metrizable spaces and $\text{card}(\mu(f)) \leq n + 1$. If $e - \dim X \leq L$, then $e - \dim Y \leq \underbrace{L \star L \star \dots \star L}_{n+1}$.*

Proof. Let $\mu(f) = \{k_1, k_2, \dots, k_\ell\}$ and $Y^i = \{y \in Y : \text{card}(f^{-1}(y)) = k_i\}$ for all $i = 1, \dots, \ell$. Obviously, $Y = \bigcup_{i=1}^{\ell} Y^i$. Since the extension dimension of a union of two subsets of any metrizable space is $\leq$ the join of their extension dimensions [9, Theorem A], it suffices to prove that $e - \dim Y^i \leq L$ for any i .

Passing to the subsets $Y^i \subset Y$ and $X^i = f^{-1}(Y^i) \subset X$, we can assume that all fibers $f^{-1}(y)$, $y \in Y$, have cardinality exactly m for some $m \geq 1$. Now, fix a metric d on X and consider the space $\exp_m(X) = \{F \subset X : \text{card}(F) \leq m\}$ equipped with the Vietoris topology, or equivalently, with the Hausdorff metric generated by d . There exists a natural continuous map $\exp_m(f): \exp_m(X) \rightarrow \exp_m(Y)$ defined by $\exp_m(f)(F) = f(F)$. Obviously, $\exp_m(f)$ is the restriction of the map $\exp_m(\beta f): \exp_m(\beta X) \rightarrow \exp_m(\beta Y)$ on $\exp_m(X)$. Since f

is perfect, the last observation implies that $\exp_m(f)$ is also perfect. So, $Z = \exp_m(f)^{-1}(Y)$ is a closed subset of $\exp_m(X)$ because Y is closed in $\exp_m(Y)$.

There is also a map $p: X^m \rightarrow \exp_m(X)$, $p((x_1, \dots, x_m)) = \{x_1, \dots, x_m\}$. It is easily seen that p is perfect, so is $p|_T$, where $T = p^{-1}(Z)$. Moreover, Z being closed in $\exp_m(X)$ yields that T is closed in X^m . Let us consider the projection $\pi: T \rightarrow X$, $\pi((x_1, \dots, x_m)) = x_1$. Since $(x_1, \dots, x_m) \in T$ if and only if $f(x_1) = \dots = f(x_m)$, we have $\pi^{-1}(K) \subset f^{-1}(f(K))^m \cap T$ for every $K \subset X$. This implies that π is finite-to-one and $\pi^{-1}(K)$ is compact provided $K \subset X$ is compact. Hence, π is a perfect light map, and by [6, Theorem 1.2], $e - \dim T \leq e - \dim X \leq L$.

Next step is to show that $e - \dim U \leq L$, where U is the subset of Z such that every $F \in U$ consists of m points. Since U is open in Z , so is $T_U = p^{-1}(U)$ in T . Consequently, by the countable sum theorem, $e - \dim T_U \leq e - \dim T \leq L$. Now, for every $\varepsilon > 0$ let

$$E_\varepsilon = \{(x_1, \dots, x_m) \in T_U : d(x_i, x_j) \geq \varepsilon \text{ for } i \neq j\}.$$

If C is a closed subset of E_ε of d_m -diameter $\leq \varepsilon/2$, where d_m is the metric on X^m defined by the same equality as in Proposition 4.1, then $p|_C$ is a homeomorphism. Indeed, suppose $p((x_1, \dots, x_m)) = p((y_1, \dots, y_m))$ for some $(x_1, \dots, x_m), (y_1, \dots, y_m) \in C$. Then $\{x_1, \dots, x_m\} = \{y_1, \dots, y_m\}$ and, since $d(x_i, y_i) \leq \varepsilon/2$ for all i , it follows that $x_i = y_i$, $i \in \{1, \dots, m\}$. So, $p|_C$ is injective. On the other hand $p: T_U \rightarrow U$ is a perfect map and C is closed in T_U because $E_\varepsilon \subset T_U$ is closed. Hence, $p|_C$ is a homeomorphism with $p(C)$ being a closed subset of U . Obviously, the sets $E_{1/k}$, $k \geq 1$, form a closed covering of T_U . For each k consider a locally finite in T_U covering Θ_k of $E_{1/k}$ with each $H \in \Theta_k$ being a closed set in T_U of d_m -diameter $\leq 1/2k$. Therefore, $\{p(H) : H \in \Theta_k\}$ is a locally finite family in U covering $p(E_{1/k})$ and consisting of closed sets in U with $e - \dim p(H) \leq L$. Consequently, $D_k = \bigcup \{p(H) : H \in \Theta_k\}$ is closed in U and $e - \dim D_k \leq L$. Finally, since $U = \bigcup_{k=1}^{\infty} D_k$, we obtain $e - \dim U \leq L$.

The last step is to show that $e - \dim Y \leq L$. We define the sets

$$Y_\varepsilon = \{y \in Y : d(x', x'') > \varepsilon \text{ for every } x' \neq x'' \in f^{-1}(y)\},$$

$\varepsilon > 0$. Using that f is a perfect map and any fibers of f contains exactly m points, one can show that each Y_ε is open in Y and all maps $g_\varepsilon: Y_\varepsilon \rightarrow U$, $g_\varepsilon(y) = f^{-1}(y)$, are embeddings. Hence, $e - \dim Y_\varepsilon \leq L$ because $e - \dim U \leq L$ yields $e - \dim A \leq L$ for every subset $A \subset U$. To complete the proof, observe that $Y = \bigcup_{k=1}^{\infty} Y_{1/k}$ with each $Y_{1/k}$ being an F_σ subset of Y implies $e - \dim Y \leq L$. $\square$

Acknowledgments. The authors would like to express their sincere thanks to E.G.Sklyarenko and A.V.Zarelua for fruitful and stimulating discussions.

REFERENCES

- [1] Borel, A., et al, *Seminar on Transformation Groups*, Annals of Mathematics Study **46**(1960).
- [2] Bredon, G.E., *Sheaf theory*, Springer, New York, 1997.
- [3] Dranishnikov, A.N., *Homology dimension theory*, Uspehi Mat. Nauk **43,4(262)** (1988), 1-55 (Russian).
- [4] Dranishnikov, A.N., *Extension of mappings into CW-complexes*, Mat. Sb. **182** (1991), 1300-1310 (Russian).
- [5] Dranishnikov, A.N., Dydak, J., *Extension dimension and extension types*, Proc. of Steklov Inst. Math. **212** (1996), 1-34.
- [6] Dranishnikov, A.N., Uspenskij, V.V., *Light maps and extensional dimension*, Topology Appl. **80** (1997), 91-99.
- [7] Dranishnikov, A.N., *Cohomological dimension theory of compact metric spaces*, arXiv:math.GN/0501523.
- [8] Dydak, J., *Extension dimension for paracompact spaces*, Topology Appl. **140**, **2-3** (2004), 227-243.
- [9] Dydak, J., *Cohomology dimension and metrizable spaces. II*, Trans. Amer. Math. Soc. **348**, **4** (1996), 1647-1661.
- [10] Dydak, J., *Cohomology dimension and metrizable spaces*, Trans. Amer. Math. Soc. **337** (1993), 219-234.
- [11] Engelking, R., *Theory of Dimensions, Finite and Infinite*, Heldermann Verlag, Lemgo, 1995.
- [12] Engelking, R., *General Topology*, Heldermann Verlag, Berlin, 1989.
- [13] Freudenthal, H., *Über dimensionerhöhende stetige Abbildungen*, Sitzber. Preuss. Akad. Wiss. **5** (1932), 34-38.
- [14] Filippov, V.V., *The behavior of dimension with respect to closed mappings*, Trudy Sem. Petrovsk. **3** (1978), 177-196 (Russian).
- [15] Godement, R., *Topologie Algébrique et Théorie der Faisceaux*, Hermann, Paris, 1958.
- [16] Huber, P.J., *Homotopical cohomology and Čech cohomology*, Math. Ann. **144** (1961), 73-76.
- [17] Hurewicz, W., *Über stetiger Bilder von Punktmengen*, Koninkl. Akad. van Wetenschappen. Proc. Sect. Sci. **30** (1927), 159-165.
- [18] Hurewicz, W., *Über dimensionerhöhende stetige Abbildungen*, J. Reine u. Angew. Math. **169**(1933), 71-78.
- [19] Keesling, J., *Mappings and dimension in general metric spaces*, Pacif. J. Math. **25** (1968), 277-288.
- [20] Kuzminov, V.I., *Homological dimension theory*, Uspehi Mat. Nauk **23** (1968), 3-49 (Russian).
- [21] Levin, M., Rubin, L.R., and Schapiro, P.J., *The Mardešć factorization theorem for extension theory and C-separation*, Proc. Amer. Math. Soc. **128**, **10** (2000), 3099-3106.
- [22] Nagami, K., *Dimension Theory*, Acad. Press, N.Y., 1970.

- [23] Pasynkov, B.A., *Factorization theorems in dimension theory*, Uspehi Mat. Nauk **36** (1981), 147–175 (Russian).
- [24] Pasynkov, B.A., *A theorem for ω -maps*, Russian Math. Surveys **39**, **5** (1984), 125–153.
- [25] Pears, A.R., *Dimension Theory of General Spaces*, Cambridge University Press, Cambridge, 1975.
- [26] Ščepin, E.V., *Topology of limit spaces of uncountable inverse spectra*, Uspehi Mat. Nauk **31**, **5** (1976), 191–226 (Russian).
- [27] Shvedov, I.A., *Dimensions and soft sheaves*, General Topology and its Relation to Modern Analysis and Algebra II, Proc. Second Prague Topological Symposium 1966, Prague 1967, 347–348 (in Russian).
- [28] Sklyarenko, E.G., *Homology and Cohomology Theories of General Spaces*, in Encyclopedia of Mathematical Sciences, **50**, Springer, Berlin, 1996, p. 118–256.
- [29] Skordev, G., *Mappings that raise dimension*, Mat. Zametki **7** (1970), 697–705 (Russian).
- [30] Skordev, G., *On a theorem of Hurewicz*, Ann. de l' Univ. de Sofia, Fac. des Math. **65** (1970/1971), 1–6 (Russian).
- [31] Skordev, G., *The resolvents of a continuous mapping*, Mat. Sb. **82** (1970), 532–550 (Russian).
- [32] Zarelua, A.V., *Finite to one mappings of topological spaces and cohomological manifolds*, Sib. Mat. J. **10** (1969), 64–92 (Russian).
- [33] Zarelua, A.V., *Cohomological structure of finite to one mappings*, Trudy Tbilis. Mat. Inst. **41** (1972), Tbilisi, 100–127 (Russian).
- [34] Zarelua, A.V., *Topological and cohomological structure of zero-dimensional mappings*, Trudy Mat. Inst. Steklova (to appear).

CENTER FOR COMPLEX SYSTEMS AND VISUALIZATION, DEPARTMENT OF COMPUTER SCIENCE AND MATHEMATICS, UNIVERSITY OF BREMEN, UNIVERSITÄT-SALLE 29, 28359 BREMEN, GERMANY

E-mail address: skordev@cvis.uni-bremen.de

DEPARTMENT OF COMPUTER SCIENCE AND MATHEMATICS, NIPISSING UNIVERSITY, 100 COLLEGE DRIVE, P.O. BOX 5002, NORTH BAY, ON, P1B 8L7, CANADA

E-mail address: veskov@nipissingu.ca